arXiv:0901.4967v2 [math.AP] 4 Feb 2009

A note on the Neumann problem

BIAGIO RICCERI

Let $\Omega \subset \mathbf{R}^n$ be a bounded domain with a boundary of class C^1, let $\alpha : \Omega \to \mathbf{R}$ be a bounded measurable function with $\inf_\Omega \alpha > 0$ and let $1 < p \leq n$. On the Sobolev space $W^{1,p}(\Omega)$, we consider the norm

$$\|u\| = \left(\int_\Omega |\nabla u(x)|^p dx + \int_\Omega \alpha(x)|u(x)|^p dx \right)^{\frac{1}{p}} .$$

We denote by $\mathcal{A}_p$ the class of all Carathéodory functions $\varphi : \Omega \times \mathbf{R} \to \mathbf{R}$ such that

$$\sup_{(x,\xi)\in\Omega\times\mathbf{R}} \frac{|\varphi(x,\xi)|}{1+|\xi|^q} < +\infty ,$$

for some q with $0 < q < \frac{pn-n+p}{n-p}$ if $p < n$ and $0 < q < +\infty$ if $p = n$. Given $\varphi \in \mathcal{A}_p$, consider the following Neumann problem

$$\begin{cases} -\mathrm{div}(|\nabla u|^{p-2}\nabla u) + \alpha(x)|u|^{p-2}u = \varphi(x,u) & \text{in } \Omega \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega \end{cases} \qquad (P_\varphi)$$

where ν denotes the outward unit normal to $\partial\Omega$.

Let us recall that a weak solution of (P_φ) is any $u \in W^{1,p}$ such that

$$\int_\Omega |\nabla u(x)|^{p-2}\nabla u(x)\nabla v(x)dx + \int_\Omega \alpha(x)|u(x)|^{p-2}u(x)v(x)dx - \int_\Omega \varphi(x,u(x))v(x)dx = 0$$

for all $v \in W^{1,p}(\Omega)$.

The functionals $\Phi, J_\varphi : W^{1,p}(\Omega) \to \mathbf{R}$ defined by

$$\Phi(u) = \frac{1}{p}\|u\|^p$$

$$J_\varphi(u) = \int_\Omega \left(\int_0^{u(x)} \varphi(x,\xi)d\xi \right) dx$$

are C^1 in $W^{1,p}(\Omega)$, with derivatives given by

$$\Phi'(u)(v) = \int_\Omega |\nabla u(x)|^{p-2}\nabla u(x)\nabla v(x)dx$$

$$J'_\varphi(u)(v) = \int_\Omega \varphi(x, u(x))v(x)dx$$

for all $u, v \in W^{1,p}(\Omega)$. Consequently, the weak solutions of problem (P_φ) are exactly the critical points in $W^{1,p}(\Omega)$ of the functional $\Phi - J_\varphi$. Moreover, J'_φ is compact, while Φ' is a homeomorphism between $W^{1,p}(\Omega)$ and its dual.

In recent years, the general abstract results established in [30], [31] (see also [35] for a revisited version) and [33] have widely been applied to study existence and multiplicity of solutions for problems like (P_φ). In this connection, we refer, for instance, to [1]-[29], [32], [34], [37].

Very recently, in [36], we obtained a new general three critical points theorem. Before recalling it, let us give the following definition.

If X is a real Banach space, we denote by $\mathcal{W}_X$ the class of all functionals $\Phi : X \to \mathbf{R}$ possessing the following property: if $\{x_n\}$ is a sequence in X converging weakly to $x \in X$ and $\liminf_{n\to\infty} \Phi(x_n) \leq \Phi(x)$, then $\{x_n\}$ has a subsequence converging strongly to x.

Also, given an operator $S : X \to X^*$, we say that S admits a continuous inverse on X^* if there exists a continuous operator $T : X^* \to X$ such that $T(S(x)) = x$ for all $x \in X$.

THEOREM A ([36], Theorem 1). - *Let X be a separable and reflexive real Banach space; $\Phi : X \to \mathbf{R}$ a coercive, sequentially weakly lower semicontinuous C^1 functional, belonging to $\mathcal{W}_X$, bounded on each bounded subset of X and whose derivative admits a continuous inverse on X^*; $J : X \to \mathbf{R}$ a C^1 functional with compact derivative. Assume that Φ has a strict local minimum x_0 with $\Phi(x_0) = J(x_0) = 0$. Finally, setting*

$$\eta = \max\left\{0, \limsup_{\|x\|\to+\infty} \frac{J(x)}{\Phi(x)}, \limsup_{x\to x_0} \frac{J(x)}{\Phi(x)}\right\},$$

$$\theta = \sup_{x\in\Phi^{-1}(]0,+\infty[)} \frac{J(x)}{\Phi(x)},$$

assume that $\eta < \theta$.

Then, for each compact interval $[a, b] \subset \left]\frac{1}{\theta}, \frac{1}{\eta}\right[$ (with the conventions $\frac{1}{0} = +\infty$, $\frac{1}{+\infty} = 0$) there exists $r > 0$ with the following property: for every $\lambda \in [a, b]$ and every C^1 functional $\Psi : X \to \mathbf{R}$ with compact derivative, there exists $\hat{\mu} > 0$ such that, for each $\mu \in [0, \hat{\mu}]$, the equation

$$\Phi'(x) = \lambda J'(x) + \mu\Psi'(x)$$

has at least three solutions whose norms are less than r.

The aim of this paper is to provide an application of Theorem A to problem (P_φ).

Our main result reads as follows:

THEOREM 1. - *Let $f \in \mathcal{A}_p$ be such that*

$$\max\left\{0, \limsup_{\xi\to 0} \frac{\sup_{x\in\Omega} F(x,\xi)}{|\xi|^p}, \limsup_{|\xi|\to+\infty} \frac{\sup_{x\in\Omega} F(x,\xi)}{|\xi|^p}\right\} <$$

$$< \frac{\inf_\Omega \alpha}{\int_\Omega \alpha(x)dx} \sup_{\xi\in\mathbf{R}\setminus\{0\}} \frac{\int_\Omega F(x,\xi)dx}{|\xi|^p} \ , \tag{1}$$

where

$$F(x,\xi) = \int_0^\xi f(x,s)ds$$

for all $(x,\xi) \in \Omega \times \mathbf{R}$.

Then, if we set

$$\gamma = \frac{\int_\Omega \alpha(x)dx}{p} \inf\left\{ \frac{|\xi|^p}{\int_\Omega F(x,\xi)dx} : \int_\Omega F(x,\xi)dx > 0 \right\}$$

and

$$\delta = \frac{\inf_\Omega \alpha}{p \max\left\{0, \limsup_{\xi\to 0} \frac{\sup_{x\in\Omega} F(x,\xi)}{|\xi|^p}, \limsup_{|\xi|\to+\infty} \frac{\sup_{x\in\Omega} F(x,\xi)}{|\xi|^p}\right\}} \ ,$$

for each compact interval $[a,b] \subset]\gamma,\delta[$ *there exists* $r > 0$ *with the following property: for every* $\lambda \in [a,b]$ *and every* $g \in \mathcal{A}_p$ *there exists* $\hat{\mu} > 0$ *such that, for each* $\mu \in [0,\hat{\mu}]$, *the problem*

$$\begin{cases} -\mathrm{div}(|\nabla u|^{p-2}\nabla u) + \alpha(x)|u|^{p-2}u = \lambda f(x,u) + \mu g(x,u) & \text{in } \Omega \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega \end{cases}$$

has at least three weak solutions whose norms in $W^{1,p}(\Omega)$ *are less than* r.

PROOF. We are going to apply Theorem A taking $X = W^{1,p}(\Omega)$, and

$$\Phi(u) = \frac{1}{p}\|u\|^p \ ,$$

$$J(u) = \int_\Omega F(x,u(x))dx$$

for all $u \in X$.

Note that, by a classical result on uniformly convex spaces, $\Phi \in \mathcal{W}_{W^{1,p}(\Omega)}$.

Also, set

$$\rho_1 = \limsup_{\xi\to 0} \frac{\sup_{x\in\Omega} F(x,\xi)}{|\xi|^p} \ ,$$

$$\rho_2 = \limsup_{|\xi|\to+\infty} \frac{\sup_{x\in\Omega} F(x,\xi)}{|\xi|^p} \ .$$

Fix $\epsilon > 0$ and choose $\delta_1, \delta_2 > 0$ so that

$$F(x,\xi) \le (\rho_1 + \epsilon)|\xi|^p \tag{2}$$

for all $(x,\xi) \in \Omega \times [-\delta_1, \delta_1]$, and

$$F(x,\xi) \le (\rho_2 + \epsilon)|\xi|^p \tag{3}$$

for all $(x,\xi) \in \Omega \times (\mathbf{R} \setminus [-\delta_2, \delta_2])$. Due to (2) and (3), since F is bounded on on each bounded subset of $\Omega \times \mathbf{R}$, we can choose $c > 0$ such that

$$F(x,\xi) \le (\rho_1 + \epsilon + c)|\xi|^p$$

for all $(x,\xi) \in \Omega \times \mathbf{R}$. So, if we fix $q > p$ (with $q \le \frac{pn-n+p}{n-p}$ if $p < n$), by continuous embedding, for some $c_1 > 0$ (independent of ϵ), one has

$$J(u) \le (\rho_1 + \epsilon) \int_\Omega |u(x)|^p dx + c_1 \|u\|^q$$

for all $u \in X$. Hence, for each $u \in X \setminus \{0\}$, we have

$$\frac{J(u)}{\|u\|^p} \le \frac{(\rho_1 + \epsilon) \int_\Omega |u(x)|^p dx}{\|u\|^p} + c_1 \|u\|^{q-p} \le \frac{\rho_1 + \epsilon}{\inf_\Omega \alpha} + c_1 \|u\|^{q-p}$$

and hence

$$\limsup_{u \to 0} \frac{J(u)}{\Phi(u)} \le \frac{p(\rho_1 + \epsilon)}{\inf_\Omega \alpha} \ . \tag{4}$$

Further, by (3) again, for each $u \in X \setminus \{0\}$, we have

$$\frac{J(u)}{\|u\|^p} = \frac{\int_{\Omega(|u| \le \delta_2)} F(x, u(x)) dx}{\|u\|^p} + \frac{\int_{\Omega(|u| > \delta_2)} F(x, u(x)) dx}{\|u\|^p} \le$$

$$\le \frac{\mathrm{meas}(\Omega) \sup_{\Omega \times [-\delta_2, \delta_2]} F}{\|u\|^p} + \frac{(\rho_2 + \epsilon) \int_\Omega |u(x)|^p dx}{\|u\|^p} \le \frac{\mathrm{meas}(\Omega) \sup_{\Omega \times [-\delta_2, \delta_2]} F}{\|u\|^p} + \frac{\rho_2 + \epsilon}{\inf_\Omega \alpha}$$

and hence

$$\limsup_{\|u\| \to +\infty} \frac{J(u)}{\Phi(u)} \le \frac{p(\rho_2 + \epsilon)}{\inf_\Omega \alpha} \ . \tag{5}$$

Since ϵ is arbitrary, (4) and (5) give

$$\max \left\{ \limsup_{\|u\| \to +\infty} \frac{J(u)}{\Phi(u)}, \limsup_{u \to 0} \frac{J(u)}{\Phi(u)} \right\} \le \frac{p}{\inf_\Omega \alpha} \max\{\rho_1, \rho_2\} \ . \tag{6}$$

Moreover, it is clear that

$$\frac{p}{\int_\Omega \alpha(x) dx} \sup_{\xi \in \mathbf{R} \setminus \{0\}} \frac{\int_\Omega F(x,\xi) dx}{|\xi|^p} \le \sup_{u \in X \setminus \{0\}} \frac{J(u)}{\Phi(u)} \ . \tag{7}$$

Then, from (1), (6) and (7), we get

$$\max\left\{0,\ \limsup_{\|u\|\to+\infty}\frac{J(u)}{\Phi(u)},\limsup_{u\to 0}\frac{J(u)}{\Phi(u)}\right\}<\sup_{u\in X\setminus\{0\}}\frac{J(u)}{\Phi(u)}\ .$$

Hence, all the assumptions of Theorem A are satisfied (with $x_0=0$). The conclusion then follows taking $\Psi=J_g$ and remarking that, due to (6) and (7), one has $[\gamma,\delta]\subseteq\left[\frac{1}{\theta},\frac{1}{\eta}\right]$. $\triangle$

We conclude pointing out an application of Theorem 1 in a case where f is a polynomial in ξ.

PROPOSITION 1. - *Let $n=3$, let $\beta:\Omega\to\mathbf{R}$ be a bounded measurable function, with $\inf_\Omega\beta\geq 0$ and $\int_\Omega\beta(x)dx>0$, and let $a,b,c\in\mathbf{R}$, with $c>0$ and $a>-\frac{2b^2}{9c}$. When $a>0$, assume that*

$$\frac{\sup_\Omega\beta\int_\Omega\alpha(x)dx}{\inf_\Omega\alpha\int_\Omega\beta(x)dx}<1+\frac{2b^2}{9ac}\ . \tag{8}$$

Set

$$\gamma_0=\frac{9c\int_\Omega\alpha(x)dx}{(9ac+2b^2)\int_\Omega\beta(x)dx}$$

and

$$\delta_0=\frac{\inf_\Omega\alpha}{\max\{0,a\}\sup_\Omega\beta}\ .$$

Then, for each compact interval $A\subset]\gamma_0,\delta_0[$, there exists $r>0$ with the following property: for every $\lambda\in A$ and every $g\in\mathcal{A}_2$ there exists $\hat\mu>0$ such that, for each $\mu\in[0,\hat\mu]$, the problem

$$\begin{cases}-\Delta u+\alpha(x)u=\lambda\beta(x)(au+bu^2-cu^3)+\mu g(x,u) & \text{in }\Omega\\ \\ \frac{\partial u}{\partial\nu}=0 & \text{on }\partial\Omega\end{cases}$$

has at least three weak solutions whose norms in $W^{1,2}(\Omega)$ are less than r.

PROOF. Set

$$f(x,\xi)=\beta(x)(a\xi+b\xi^2-c\xi^3)$$

for all $(x,\xi)\in\Omega\times\mathbf{R}$. Note that $f\in\mathcal{A}_2$ as $n=3$. Moreover, if we set

$$\Gamma=\left\{\xi\in\mathbf{R}:\frac{a}{2}+\frac{b}{3}\xi-\frac{c}{4}\xi^2\geq 0\right\}$$

we clearly have

$$\frac{\sup_{x\in\Omega}F(x,\xi)}{\xi^2}=\begin{cases}\sup_\Omega\beta\left(\frac{a}{2}+\frac{b}{3}\xi-\frac{c}{4}\xi^2\right) & \text{if }\xi\in\Gamma\\ \\ \inf_\Omega\beta\left(\frac{a}{2}+\frac{b}{3}\xi-\frac{c}{4}\xi^2\right) & \text{if }\xi\in\mathbf{R}\setminus\Gamma\ .\end{cases}$$

From this, we obtain

$$\lim_{|\xi|\to+\infty} \frac{\sup_{x\in\Omega} F(x,\xi)}{\xi^2} \leq 0$$

(since $c > 0$ and $\inf_\Omega \beta \geq 0$) as well as

$$\lim_{\xi\to 0} \frac{\sup_{x\in\Omega} F(x,\xi)}{\xi^2} = \begin{cases} \frac{a \sup_\Omega \beta}{2} & \text{if } a > 0 \\ \frac{a \inf_\Omega \beta}{2} & \text{if } a \leq 0 \ . \end{cases}$$

Consequently, we have

$$\max\left\{0, \lim_{\xi\to 0} \frac{\sup_{x\in\Omega} F(x,\xi)}{\xi^2}, \lim_{|\xi|\to+\infty} \frac{\sup_{x\in\Omega} F(x,\xi)}{\xi^2}\right\} = \frac{\max\{0,a\} \sup_\Omega \beta}{2} \ . \tag{9}$$

On the other hand, we also have

$$\sup_{\xi\in\mathbf{R}\setminus\{0\}} \frac{\int_\Omega F(x,\xi)dx}{\xi^2} = \left(\frac{a}{2} + \frac{b^2}{9c}\right) \int_\Omega \beta(x)dx \ . \tag{10}$$

At this point, the validity of (1) (for $p = 2$) follows from (8), (9) and (10) when $a > 0$. When $a \leq 0$, (1) follows from (9) and (10) as the right-hand side of (10) is positive, by assumption. So, f satisfies the assumptions of Theorem 1, with $p = 2$. Now, the conclusion follows as $\gamma_0 = \gamma$ and $\delta_0 = \delta$. △

References

[1] G. ANELLO and G. CORDARO, *Existence of solutions of the Neumann problem involving the p-Laplacian via a variational principle of Ricceri*, Arch. Math. (Basel), **79** (2002), 274-287.

[2] G. ANELLO and G. CORDARO, *An existence theorem for the Neumann problem involving the p-Laplacian*, J. Convex Anal., **10** (2003), 185-198.

[3] G. ANELLO and G. CORDARO, *Infinitely many positive solutions for the Neumann problem involving the p-Laplacian*, Colloq. Math., **97** (2003), 221-231.

[4] G. ANELLO, *Existence of infinitely many weak solutions for a Neumann problem*, Nonlinear Anal., **57** (2004), 199-209.

[5] G. ANELLO, *A multiplicity theorem for critical points of functionals on reflexive Banach spaces*, Arch. Math. (Basel), **82** (2004), 172-179.

[6] G. ANELLO, *Existence and multiplicity of solutions to a perturbed Neumann problem*, Math. Nachr., **280** (2007), 1755-1764.

[7] G. ANELLO, *Remarks on a multiplicity theorem for a perturbed Neumann problem*, J. Math. Anal. Appl., **346** (2008), 274-279.

[8] D. AVERNA and G. BONANNO, *Three solutions for a Neumann boundary value problem involving the p-Laplacian*, Matematiche, **60** (2005), 81-91.

[9] G. BILOTTA, *Existence of infinitely many solutions for a quasilinear Neumann problem*, Panamer. Math. J., **13** (2003), 1-18.

[10] G. BONANNO, *Multiple solutions for a Neumann boundary value problem*, J. Nonlinear Convex Anal., **4** (2003), 287-290.

[11] G. BONANNO, *Some remarks on a three critical points theorem*, Nonlinear Anal., **54** (2003), 651-665.

[12] G. BONANNO and P. CANDITO, *Three solutions to a Neumann problem for elliptic equations involving the p-Laplacian*, Arch. Math. (Basel), **80** (2003), 424-429.

[13] F. CAMMAROTO, A. CHINNÌ and B. DI BELLA, *Some multiplicity results for quasilinear Neumann problems*, Arch. Math. (Basel), **86** (2006), 154-162.

[14] P. CANDITO, *Infinitely many solutions to the Neumann problem for elliptic equations involving the p-Laplacian and with discontinuous nonlinearities*, Proc. Edinb. Math. Soc., **45** (2002), 397-409.

[15] G. CORDARO, *Multiple solutions to a perturbed Neumann problem*, Studia Math., **178** (2007), 167-175.

[16] G. DAI, *Infinitely many solutions for a Neumann-type differential inclusion problem involving the $p(x)$-Laplacian*, Nonlinear Anal., to appear.

[17] A. G. DI FALCO, *Infinitely many solutions of the Neumann problem for quasilinear elliptic systems*, Matematiche, **58** (2003), 117-130.

[18] S. EL MANOUNI and M. KBIRI ALAOUI, *A result on elliptic systems with Neumann conditions via Ricceri's three critical point theorem*, Nonlinear Anal., to appear.

[19] X. L. FAN and S. G. DENG, *Remarks on Ricceri's variational principle and applications to $p(x)$-Laplacian equations*, Nonlinear Anal., **67** (2007), 3064-3075.

[20] X. L. FAN and C. JI, *Existence of infinitely many solutions for a Neumann problem involving the $p(x)$-Laplacian*, J. Math. Anal. Appl., **334** (2007), 248-260.

[21] F. FARACI, *Multiplicity results for a Neumann problem involving the p-Laplacian*, J. Math. Anal. Appl., **277** (2003), 180-189.

[22] F. FARACI and A. KRISTÁLY, *On an open question of Ricceri concerning a Neumann problem*, Glasg. Math. J., **49** (2007), 189-195.

[23] A. IANNIZZOTTO, *A sharp existence and localization theorem for a Neumann problem*, Arch. Math. (Basel), **82** (2004), 352-360.

[24] A. KRISTÁLY, M. MIHĂILESCU and V. RĂDULESCU, *Two non-trivial solutions for a non-homogeneous Neumann problem: an Orlicz-Sobolev setting*, Proc. Royal Soc. Edinburgh Sect. A, to appear.

[25] A. KRISTÁLY, N. S. PAPAGEORGIOU and Cs. VARGA, *Multiple solutions for a class of Neumann elliptic problems on compact Riemannian manifolds with boundary*, Canad. Math. Bull., to appear.

[26] Q. LIU, *Existence of three solutions for $p(x)$-Laplacian equations*, Nonlinear Anal., **68** (2008), 2119-2127.

[27] S. A. MARANO and D. MOTREANU, *Infinitely many critical points of non-differentiable functions and applications to a Neumann type problem involving the p-Laplacian*, J. Differential Equations, **182** (2002), 108-120.

[28] M. MIHĂILESCU, *Existence and multiplicity of solutions for a Neumann problem involving the $p(x)$-Laplace operator*, Nonlinear Anal., **67** (2007), 1419-1425.

[29] D. S. MOSCHETTO, *A quasilinear Neumann problem involving the $p(x)$-Laplacian*, Nonlinear Anal., to appear.

[30] B. RICCERI, *A general variational principle and some of its applications*, J. Comput. Appl. Math., **113** (2000), 401-410.

[31] B. RICCERI, *On a three critical points theorem*, Arch. Math. (Basel), **75** (2000), 220-226.

[32] B. RICCERI, *Infinitely many solutions of the Neumann problem for elliptic equations involving the p-Laplacian*, Bull. London Math. Soc., **33** (2001), 331-340.

[33] B. RICCERI, *Sublevel sets and global minima of coercive functionals and local minima of their perturbations*, J. Nonlinear Convex Anal., **5** (2004), 157-168.

[34] B. RICCERI, *A multiplicity theorem for the Neumann problem*, Proc. Amer. Math. Soc., **134** (2006), 1117-1124.

[35] B. RICCERI, *A three critical points theorem revisited*, Nonlinear Anal., to appear.

[36] B. RICCERI, *A further three critical points theorem*, preprint.

[37] X. SHI and X. DING, *Existence and multiplicity of solutions for a general $p(x)$-Laplacian Neumann problem*, Nonlinear Anal., to appear.

Department of Mathematics
University of Catania
Viale A. Doria 6
95125 Catania
Italy
e-mail address: ricceri@dmi.unict.it